



## New Criterion for the Riemann Hypothesis

---

Frank Vega

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

November 29, 2023

# New Criterion for the Riemann Hypothesis

Frank Vega

*To my mother*

**Abstract.** The Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part  $\frac{1}{2}$ . It is considered by many to be the most important unsolved problem in pure mathematics. Let  $\Psi(n) = n \cdot \prod_{q|n} \left(1 + \frac{1}{q}\right)$  denote the Dedekind  $\Psi$  function where  $q | n$  means the prime  $q$  divides  $n$ . Define, for  $n \geq 3$ ; the ratio  $R(n) = \frac{\Psi(n)}{n \cdot \log \log n}$  where  $\log$  is the natural logarithm. Let  $N_n = 2 \cdot \dots \cdot q_n$  be the primorial of order  $n$ . There are several statements equivalent to the Riemann hypothesis. We state that if for each large enough prime number  $q_n$ , there exists another prime  $q_{n'} > q_n$  such that  $R(N_{n'}) \leq R(N_n)$ , then the Riemann hypothesis is true. In this note, using our criterion, we prove that the Riemann hypothesis is true.

**Mathematics Subject Classification (2010).** Primary 11M26; Secondary 11A41.

**Keywords.** Elementary Number Theory, Riemann hypothesis, prime numbers, Riemann zeta function, Chebyshev function.

## 1. Introduction

The Riemann hypothesis was proposed by Bernhard Riemann (1859). The Riemann hypothesis belongs to the Hilbert's eighth problem on Hilbert's list of twenty-three unsolved problems. This is one of the Clay Mathematics Institute's Millennium Prize Problems. In mathematics, the Chebyshev function  $\theta(x)$  is given by

$$\theta(x) = \sum_{q \leq x} \log q$$

with the sum extending over all prime numbers  $q$  that are less than or equal to  $x$ , where  $\log$  is the natural logarithm. Leonhard Euler studied the following value of the Riemann zeta function (1734) [1].

**Proposition 1.1.** We define [1, (1) pp. 1070]:

$$\zeta(2) = \prod_{k=1}^{\infty} \frac{q_k^2}{q_k^2 - 1} = \frac{\pi^2}{6},$$

where  $q_k$  is the  $k$ th prime number (We also use the notation  $q_n$  to denote the  $n$ th prime number). By definition, we have

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1}{n^2},$$

where  $n$  denotes a natural number. Leonhard Euler proved in his solution to the Basel problem that

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \prod_{k=1}^{\infty} \frac{q_k^2}{q_k^2 - 1} = \frac{\pi^2}{6},$$

where  $\pi \approx 3.14159$  is a well-known constant linked to several areas in mathematics such as number theory, geometry, etc.

The number  $\gamma \approx 0.57721$  is the Euler-Mascheroni constant which is defined as

$$\begin{aligned} \gamma &= \lim_{n \rightarrow \infty} \left( -\log n + \sum_{k=1}^n \frac{1}{k} \right) \\ &= \int_1^{\infty} \left( -\frac{1}{x} + \frac{1}{[x]} \right) dx. \end{aligned}$$

Here,  $[\dots]$  represents the floor function. In number theory,  $\Psi(n) = n \cdot \prod_{q|n} \left( 1 + \frac{1}{q} \right)$  is called the Dedekind  $\Psi$  function, where  $q | n$  means the prime  $q$  divides  $n$ . We say that  $\text{Dedekind}(q_n)$  holds provided that

$$\prod_{q \leq q_n} \left( 1 + \frac{1}{q} \right) > \frac{e^\gamma}{\zeta(2)} \cdot \log \theta(q_n).$$

Next, we have the Solé and Planat Theorem:

**Proposition 1.2.**  $\text{Dedekind}(q_n)$  holds for all prime numbers  $q_n > 3$  if and only if the Riemann hypothesis is true [8, Theorem 4.2 pp. 5].

A natural number  $N_n$  is called a primorial number of order  $n$  precisely when,

$$N_n = \prod_{k=1}^n q_k.$$

We define  $R(n) = \frac{\Psi(n)}{n \cdot \log \log n}$  for  $n \geq 3$ .  $\text{Dedekind}(q_n)$  holds if and only if  $R(N_n) > \frac{e^\gamma}{\zeta(2)}$  is satisfied.

**Proposition 1.3.** Unconditionally on Riemann hypothesis, we know that [8, Proposition 3. pp. 3]:

$$\lim_{n \rightarrow \infty} R(N_n) = \frac{e^\gamma}{\zeta(2)}.$$

**Proposition 1.4.** *For all prime numbers  $q_n > 5$  [3, Theorem 1.1. pp. 358]:*

$$\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right) < e^\gamma \cdot \log \theta(q_n).$$

The well-known asymptotic notation  $\Omega$  was introduced by Godfrey Harold Hardy and John Edensor Littlewood [4]. In 1916, they also introduced the two symbols  $\Omega_R$  and  $\Omega_L$  defined as [5]:

$$f(x) = \Omega_R(g(x)) \text{ as } x \rightarrow \infty \text{ if } \limsup_{x \rightarrow \infty} \frac{f(x)}{g(x)} > 0;$$

$$f(x) = \Omega_L(g(x)) \text{ as } x \rightarrow \infty \text{ if } \liminf_{x \rightarrow \infty} \frac{f(x)}{g(x)} < 0.$$

After that, many mathematicians started using these notations in their works. From the last century, these notations  $\Omega_R$  and  $\Omega_L$  changed as  $\Omega_+$  and  $\Omega_-$ , respectively. There is another notation:  $f(x) = \Omega_\pm(g(x))$  (meaning that  $f(x) = \Omega_+(g(x))$  and  $f(x) = \Omega_-(g(x))$  are both satisfied). Nowadays, the notation  $f(x) = \Omega_+(g(x))$  has survived and it is still used in analytic number theory as [9]:

$$f(x) = \Omega_+(g(x)) \text{ if } \exists k > 0 \forall x_0 \exists x > x_0: f(x) \geq k \cdot g(x)$$

which has the same meaning to the Hardy and Littlewood older notation. For  $x \geq 2$ , the function  $f$  was introduced by Nicolas in his seminal paper as [7, Theorem 3 pp. 376], [2, (5.5) pp. 111]:

$$f(x) = e^\gamma \cdot \log \theta(x) \cdot \prod_{q \leq x} \left(1 - \frac{1}{q}\right).$$

Finally, we have the Nicolas Theorem:

**Proposition 1.5.** *If the Riemann hypothesis is false then there exists a real  $b$  with  $0 < b < \frac{1}{2}$  such that, as  $x \rightarrow \infty$  [7, Theorem 3 (c) pp. 376], [2, Theorem 5.29 pp. 131]:*

$$\log f(x) = \Omega_\pm(x^{-b}).$$

Besides, we based our approach on the following property:

**Proposition 1.6.** *For  $x > -1$  and  $r > 1$  [6, pp. 1]:*

$$(1 + x)^r \geq 1 + r \cdot x.$$

Putting all together yields a proof for the Riemann hypothesis.

## 2. Central Lemma

Several analogues of the Riemann hypothesis have already been proved. Many authors expect (or at least hope) that it is true. However, there exist some implications in case of the Riemann hypothesis could be false. The following is a key Lemma.

**Lemma 2.1.** *If the Riemann hypothesis is false, then there exist infinitely many prime numbers  $q_n$  such that  $\text{Dedekind}(q_n)$  fails (i.e.  $\text{Dedekind}(q_n)$  does not hold).*

*Proof.* The function  $g$  is defined as [8, Theorem 4.2 pp. 5]:

$$g(x) = \frac{e^\gamma}{\zeta(2)} \cdot \log \theta(x) \cdot \prod_{q \leq x} \left(1 + \frac{1}{q}\right)^{-1}.$$

By Proposition 1.2, the Riemann hypothesis is false whenever there exists some natural number  $x_0 \geq 5$  such that  $g(x_0) > 1$  or equivalent  $\log g(x_0) > 0$ . It was proven the following bound [8, Theorem 4.2 pp. 5]:

$$\log g(x) \geq \log f(x) - \frac{2}{x}.$$

By Proposition 1.5, if the Riemann hypothesis is false, then there is a real number  $0 < b < \frac{1}{2}$  such that there exist infinitely many natural numbers  $x$  for which  $\log f(x) = \Omega_+(x^{-b})$ . Actually Nicolas proved that  $\log f(x) = \Omega_\pm(x^{-b})$ , but we only need to use the notation  $\Omega_+$  under the domain of the natural numbers. According to the Hardy and Littlewood definition, this would mean that

$$\exists k > 0, \forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} (y > y_0): \log f(y) \geq k \cdot y^{-b}.$$

The previous inequality is also  $\log f(y) \geq (k \cdot y^{-b} \cdot \sqrt{y}) \cdot \frac{1}{\sqrt{y}}$ , but we notice that

$$\lim_{y \rightarrow \infty} (k \cdot y^{-b} \cdot \sqrt{y}) = \infty$$

for every possible values of  $k > 0$  and  $0 < b < \frac{1}{2}$ . Now, this implies that

$$\forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} (y > y_0): \log f(y) \geq \frac{1}{\sqrt{y}}.$$

Note that, the value of  $k$  is not necessary in our previous expression. In this way, if the Riemann hypothesis is false, then there exist infinitely many natural numbers  $x$  such that  $\log f(x) \geq \frac{1}{\sqrt{x}}$ . Since  $\frac{1}{\sqrt{x_0}} > \frac{2}{x_0}$  for  $x_0 \geq 5$ , then it would be infinitely many natural numbers  $x_0$  such that  $\log g(x_0) > 0$ . In addition, if  $\log g(x_0) > 0$  for some natural number  $x_0 \geq 5$ , then  $\log g(x_0) = \log g(q_n)$  where  $q_n$  is the greatest prime number such that  $q_n \leq x_0$ . Actually, we know that

$$\prod_{q \leq x_0} \left(1 + \frac{1}{q}\right)^{-1} = \prod_{q \leq q_n} \left(1 + \frac{1}{q}\right)^{-1}$$

and

$$\theta(x_0) = \theta(q_n)$$

according to the definition of the Chebyshev function. □

### 3. Main Insight

This is the main insight.

**Lemma 3.1.** *The Riemann hypothesis is true whenever for each large enough prime number  $q_n$ , there exists another prime  $q_{n'} > q_n$  such that*

$$R(N_{n'}) \leq R(N_n).$$

*Proof.* By Lemma 2.1, if the Riemann hypothesis is false and the inequality

$$R(N_{n'}) \leq R(N_n)$$

is satisfied for each large enough prime number  $q_n$ , then there exists an infinite subsequence of natural numbers  $n_i$  such that

$$R(N_{n_{i+1}}) \leq R(N_{n_i}),$$

$q_{n_{i+1}} > q_{n_i}$  and  $\text{Dedekind}(q_{n_i})$  fails. By Proposition 1.3, this is a contradiction with the fact that

$$\liminf_{n \rightarrow \infty} R(N_n) = \lim_{n \rightarrow \infty} R(N_n) = \frac{e^\gamma}{\zeta(2)}.$$

By definition of the limit inferior for any positive real number  $\varepsilon$ , only a finite number of elements of  $R(N_n)$  are less than  $\frac{e^\gamma}{\zeta(2)} - \varepsilon$ . This contradicts the existence of such previous infinite subsequence and thus, the Riemann hypothesis must be true.  $\square$

### 4. Main Theorem

This is the main theorem.

**Theorem 4.1.** *The Riemann hypothesis is true.*

*Proof.* By Lemma 3.1, the Riemann hypothesis is true whenever

$$R(N_{n'}) \leq R(N_n)$$

is satisfied for large enough prime numbers  $q_{n'} > q_n$ . That is the same as

$$\frac{\prod_{q \leq q_{n'}} \left(1 + \frac{1}{q}\right)}{\log \theta(q_{n'})} \leq \frac{\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right)}{\log \theta(q_n)}$$

and

$$\frac{\prod_{q \leq q_{n'}} \left(1 + \frac{1}{q}\right)}{\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right)} \leq \frac{\log \theta(q_{n'})}{\log \theta(q_n)}$$

which is

$$\log \log \theta(q_{n'}) - \log \log \theta(q_n) \geq \sum_{q_n < q \leq q_{n'}} \log \left(1 + \frac{1}{q}\right)$$

after of applying the logarithm to the both sides and distribute the terms. That is equivalent to

$$1 - \frac{\log \log \theta(q_n)}{\log \log \theta(q_{n'})} \geq \frac{\sum_{q_n < q \leq q_{n'}} \log \left(1 + \frac{1}{q}\right)}{\log \log \theta(q_{n'})}$$

after dividing both sides by  $\log \log \theta(q_{n'})$ . We can apply the exponentiation to the both sides in order to obtain that

$$\exp \left(1 - \frac{\log \log \theta(q_n)}{\log \log \theta(q_{n'})}\right) \geq \left( \prod_{q_n < q \leq q_{n'}} \log \left(1 + \frac{1}{q}\right) \right)^{\frac{1}{\log \log \theta(q_{n'})}}.$$

We can take an arbitrary large enough prime  $q_{n'}$  such that

$$\frac{\log \log \theta(q_n)}{\log \log \theta(q_{n'})} \approx 0.$$

Certainly, we could have

$$\exp \left(1 - \frac{\log \log \theta(q_n)}{\log \log \theta(q_{n'})}\right) \approx e$$

for an arbitrary prime number  $q_{n'}$  much greater than  $q_n$ . Consequently, we are able to state that

$$e \cdot \alpha \geq \left( \prod_{q_n < q \leq q_{n'}} \log \left(1 + \frac{1}{q}\right) \right)^{\frac{1}{\log \log \theta(q_{n'})}}$$

for a positive value  $1 > \alpha \geq \left(1 - \frac{1}{\log \theta(q_{n'})}\right)$  such that

$$\exp \left(1 - \frac{\log \log \theta(q_n)}{\log \log \theta(q_{n'})}\right) \geq e \cdot \alpha.$$

For large enough prime  $q_{n'}$ , we have

$$e = (\log \theta(q_{n'}))^{\frac{1}{\log \log \theta(q_{n'})}}$$

since  $e = x^{\frac{1}{\log x}}$  for  $x > 0$ . So, we deduce that

$$\alpha^{\log \log \theta(q_{n'})} \cdot \log \theta(q_{n'}) \geq \prod_{q_n < q \leq q_{n'}} \log \left(1 + \frac{1}{q}\right).$$

That is equal to

$$\alpha^{\log \log \theta(q_{n'})} \cdot e^\gamma \cdot \log \theta(q_{n'}) \geq e^\gamma \cdot \prod_{q_n < q \leq q_{n'}} \log \left(1 + \frac{1}{q}\right).$$

By Proposition 1.6, we can see that

$$\begin{aligned} \alpha^{\log \log \theta(q_{n'})} &\geq \left(1 - \frac{1}{\log \theta(q_{n'})}\right)^{\log \log \theta(q_{n'})} \\ &\geq \left(1 - \frac{\log \log \theta(q_{n'})}{\log \theta(q_{n'})}\right) \end{aligned}$$

for large enough prime  $q_{n'}$ . By Proposition 1.4, we know that

$$e^\gamma \cdot \log \theta(q_{n'}) > \prod_{q \leq q_{n'}} \log \left( 1 + \frac{1}{q} \right).$$

Hence, it is enough to show that

$$\left( 1 - \frac{\log \log \theta(q_{n'})}{\log \theta(q_{n'})} \right) \geq e^\gamma \cdot \prod_{q \leq q_n} \log \left( 1 + \frac{1}{q} \right)^{-1}$$

which is trivially true since

$$\lim_{n' \rightarrow \infty} \left( 1 - \frac{\log \log \theta(q_{n'})}{\log \theta(q_{n'})} \right) = 1$$

and

$$\lim_{n \rightarrow \infty} \left( e^\gamma \cdot \prod_{q \leq q_n} \log \left( 1 + \frac{1}{q} \right)^{-1} \right) = 0. \quad \square$$

## 5. Conclusions

Practical uses of the Riemann hypothesis include many propositions that are considered to be true under the assumption of the Riemann hypothesis and some of them that can be shown to be equivalent to the Riemann hypothesis. Indeed, the Riemann hypothesis is closely related to various mathematical topics such as the distribution of primes, the growth of arithmetic functions, the Lindelöf hypothesis, the Large Prime Gap Conjecture, etc. A proof of the Riemann hypothesis could spur considerable advances in many mathematical areas, such as number theory and pure mathematics in general.

### Acknowledgment

Many thanks to Patrick Solé and Michel Planat for their support.

## References

- [1] Ayoub, R.: Euler and the Zeta Function. *The American Mathematical Monthly* **81**(10), 1067–1086 (1974). <https://doi.org/10.2307/2319041>
- [2] Broughan, K.: Euler's Totient Function, *Encyclopedia of Mathematics and its Applications*, vol. 1, pp. 94–143. Cambridge University Press (2017). <https://doi.org/10.1017/9781108178228.007>
- [3] Choie, Y., Lichiardopol, N., Moree, P., Solé, P.: On Robin's criterion for the Riemann hypothesis. *Journal de Théorie des Nombres de Bordeaux* **19**(2), 357–372 (2007). <https://doi.org/10.5802/jtnb.591>
- [4] Hardy, G.H., Littlewood, J.E.: Some problems of diophantine approximation: Part II. The trigonometrical series associated with the elliptic  $\vartheta$ -functions. *Acta mathematica* **37**(1), 193–239 (1914)



- [5] Hardy, G.H., Littlewood, J.E.: Contributions to the theory of the Riemann zeta-function and the theory of the distribution of primes. *Acta Mathematica* **41**, 119–196 (1916)
- [6] Kozma, L.: Useful Inequalities. Kozma’s Homepage, Useful inequalities cheat sheet. [http://www.lkozma.net/inequalities\\_cheat\\_sheet/ineq.pdf](http://www.lkozma.net/inequalities_cheat_sheet/ineq.pdf) (2011), Accessed 29 November 2023
- [7] Nicolas, J.L.: Petites valeurs de la fonction d’Euler. *Journal of Number Theory* **17**(3), 375–388 (1983). [https://doi.org/10.1016/0022-314X\(83\)90055-0](https://doi.org/10.1016/0022-314X(83)90055-0)
- [8] Solé, P., Planat, M.: Extreme values of the Dedekind  $\psi$  function. *Journal of Combinatorics and Number Theory* **3**(1), 33–38 (2011)
- [9] Tenenbaum, G.: *Introduction to Analytic and Probabilistic Number Theory*, vol. 163. American Mathematical Soc. (2015)

Frank Vega  
NataSquad  
10 rue de la Paix  
FR 75002 Paris  
France  
e-mail: [vega.frank@gmail.com](mailto:vega.frank@gmail.com)

Submitted: November 29, 2023.